



Mitochondrial DNA and Population Size

Oliver F. Berry, *et al.*

Science **314**, 1388 (2006);

DOI: 10.1126/science.314.5804.1388

***The following resources related to this article are available online at
www.sciencemag.org (this information is current as of December 1, 2006):***

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<https://www.sciencemag.org/cgi/content/full/314/5804/1388>

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<https://www.sciencemag.org/help/about/permissions.dtl>

Looking below
eusociality

1391




Parenting natural
phenomena

1393



Turning pattern of
hair growth

1397



LETTERS | BOOKS | POLICY FORUM | EDUCATION FORUM | PERSPECTIVES

LETTERS

edited by Etta Kavanagh

Balancing Communication and Safety

DEMOCRATIC SOCIETIES ARE NOW ENGAGED AGAINST TERRORIST ACTIVITIES. In such an environment, there is tension between the desire to withhold scientific information from those who would use it for ill and the need to not stifle fundamental research in the life sciences or the open communication of results. To inhibit the

“Freedom for research and communication is more necessary than ever...”

—Ehrlich

pursuit of science may suggest safety from those prepared to use science for harmful purposes, but any sense of security is false. Freedom for research and communication is more necessary than ever, and the best defense against those who would employ science as a weapon is scientific excellence. There will inevitably be worldwide communication of the results of scientific studies, but open communication is vital to peer review and an independent evaluation of research, including oversight by the executive and legislative branches

of government as well as the public. Open communication is also essential for public-health and public-safety planning, for the robust growth of business and technology, and for research that will be beneficial for society. Such openness is additionally necessary for the development of countermeasures against sinister applications of science. Preventing publication, even if that could be accomplished, will not prevent the misuse of science because sanctions will not deter those who have a malevolent intent. Secrecy instead poses the danger of enforced ignorance.

The life-sciences community has generally garnered public trust. To ensure the continued success of the scientific enterprise, it is critical to maintain and further that trust against the possibility of public misunderstanding, particularly in an ever-changing scientific and political environment. To preserve their credibility, members of the scientific community must remain sensitive to the potential that information could be misused by individuals and communities to endanger public safety and health or otherwise jeopardize national security; continuing education and responsible engagement in the wider body politic are required.

Life scientists enjoy a virtually unrestricted exchange of information; shared information has been a safeguard and a cornerstone. But legitimate threats to our national security necessitate that there be appropriate oversight of scientific research and publication. Restraints of the kind set forth by President Reagan in National Security Decision Directive (NSDD) 189 (1) are fit. However, perfect regulation is impossible because it assumes perfect compliance. While the scientific community continues to accept responsibility for principled research and communication, and regulation as a management tool, the public and the government must recognize that true national security requires scientific accomplishment and that scientific excellence requires the open communication of research and results. **SUSAN A. EHRLICH***

Judge, Arizona Court of Appeals, 1501 West Washington Street, Phoenix, AZ 85007, USA, and a member of the National Science Advisory Board on Biosecurity.

*The views expressed are the author's alone.

Reference

1. Available at www.fas.org/irp/offdocs/nsdd/nsdd-189.htm.

Glossing Over the Complexity of Water

ALTHOUGH WE APPLAUD THE RECOGNITION GIVEN BY *Science* to Freshwater Resources, the recent Special Section (25 Aug., pp. 1067–1090) missed an opportunity to highlight the multifaceted nature of water resources research. Framing “the” water problem as a search to quench a universal thirst (“A thirsty world”) glosses over critical differences in the causes of, and thus the solutions to, water problems across regions. It forces the discussion into the domains of supply augmentation and engineering and marginalizes underlying drivers of “thirst” such as rapid urbanization, economic transitions, geopolitical factors, or poverty.

Lack of access to water in many African countries, for example, is less the outcome of a first-order water scarcity than of a second-order scarcity of social resources (1). As the News story “Running out of water—and time” (J. Bohannon, p. 1085) suggests, Gaza suffers at least as much from geopolitical factors that inhibit access to money and nearby water as from the “environmental problem” of “running out of water.” Water transfers or desalination help overcome local/regional scarcity, but with important environmental, social, and economic costs (“Going against the flow,” R. Stone, H. Jia, News Focus, p. 1034; “Desalination freshens up,” R. F. Service, News, p. 1088). For example, Israel’s water management is becoming “sustainable” (“Seeking sustainability: Israel’s evolving water management strategy,” A. Tal, Perspective, p. 1081) only from a narrow technical perspective that treats as exogenous the growth in its arid south and neglects the environmental and third-order impacts of overexploiting the Jordan River. First-order scarcity metrics (“Global hydrological cycles and world water resources,” T. Oki, S. Kanae, Review, p. 1068), especially global ones, overlook such specificities and are of limited policy use.

The interdisciplinary water research community has shifted its attention to context-specific and proactive approaches such as watershed management, ecological engineering, demand management, reallocation, and collaborative/adaptive planning (2). We understand that the Special Section was not meant to be an exhaustive review of freshwater issues. But institutional, political, and economic options deserve more than cursory mention in *Science*, since it is primarily these, rather than technical fixes alone, that “offer a measure of hope for the future” (“A thirsty world,” J. Yeston *et al.*, p. 1067).

GIORGOS KALLIS, MICHAEL KIPARSKY,
ANITA MILMAN, ISHA RAY

Energy and Resources Group (ERG), University of California at Berkeley, Berkeley, CA 94720–3050, USA. E-mail: gkallis@berkeley.edu

References

1. J. Lundqvist, M. Falkenmark, C. Folke, L. Gordon, L. Ohlsson, *New Dimensions in Water Security* (FAO AGU/MISC/25/2000, UN Food and Agriculture Organization, Rome, 2000).
2. P. H. Gleick, *Water Int.* **25**, 127 (2000).
3. The Letter was written with input from the ERG Water Group.

Mitochondrial DNA and Population Size

IN THEIR REPORT “POPULATION SIZE DOES NOT influence mitochondrial genetic diversity in animals” (28 Apr., p. 570), E. Bazin *et al.* present compelling evidence that selective sweeps occur in animal mitochondrial DNA (mtDNA) and reduce genetic diversity below the level expected at mutation-drift equilibrium in some taxa. They also assert that this evidence implies that mtDNA has limited relevance to biodiversity and conservation studies. I contest this claim on two fronts.

First, the selective sweeps that they detect occur at very deep phylogenetic levels (phyla to class), which translate into deep evolution-

ary time (hundreds of millions of years). It is rare that conservation biologists are interested in how mtDNA diversity is distributed at such a level. Rather, it is standard practice that genetic diversity is interpreted in the context of a relevant, almost always closely related, control group (1). This practice is designed to account as best as possible for the potentially confounding historical, demographic, mutational, and selective variables that influence genetic diversity.

Second, it is well established that the geographical distribution of mtDNA diversity as determined by lineage-sorting, and not just diversity per se, is informative with respect to biodiversity conservation (2–4). Use of this criterion is recognized to address the very differences in accumulation or maintenance of genetic diversity within different taxa described by Bazin *et al.*—otherwise known as the “how much divergence is enough” question (3).

Clearly, conservation biologists should not ignore selective sweeps; they do occur, and sometimes rapidly (5). However, mtDNA diversity is abundant at the population, species, and genus level of animals (2), and it is here that it can be, and is, most relevant and rou-

tinely exploited for conservation purposes. This would not be the case if selective sweeps were as dominant a force as implied by Bazin *et al.* Despite their claims, Bazin *et al.*'s results have limited relevance to most standard applications of mtDNA in conservation.

OLIVER F. BERRY

School of Animal Biology, University of Western Australia, Stirling Highway, Crawley, Western Australia 6009, Australia.

References


1. J. L. Bouzat, *Genetica* **110**, 109 (2000).
2. J. C. Avise, *Phylogeography: The History and Formation of Species* (Harvard Univ. Press, Cambridge, MA, 2000).
3. C. Moritz, *Trends Ecol. Evol.* **9**, 373 (1994).
4. P. D. N. Hebert, A. Cywinska, S. L. Ball, J. R. deWaard, *Proc. R. Soc. London B* **270**, 313 (2002).
5. M. Turelli, A. A. Hoffmann, *Nature* **353**, 440 (1991).

IN A META-ANALYSIS OF GENETIC POLYMORPHISM, E. Bazin *et al.* suggest that mitochondrial DNA (mtDNA) is more profoundly affected by nonneutral evolution than nuclear loci (“Population size does not influence mitochondrial genetic diversity in animals,” Reports, 28 Apr., p. 570). This interpretation has already led some to conclude that mtDNA is of little utility in studies of evolution and conservation. It is well known

that multiple evolutionary processes must be considered in interpreting patterns of genetic diversity at any gene region (1, 2). However, dismissing mtDNA as a more biased analytical tool is neither necessary nor justified (3).

First, it is inappropriate to approximate effective population size (N_e) from census size, as is implied by Bazin *et al.*'s "intuitive" predictions. Bottlenecks, fluctuating population size, reproductive strategies, and geographic structure, none of which can be inferred reliably from present census size, profoundly impact N_e and genetic diversity (4). Indeed, invertebrate taxa and fish generally have greater census size than tetrapods, but there is also greater diversity in life history and reproductive strategies, traits that alter patterns of sequence divergence within and among taxa.

Second, the neutrality index (NI) may be inappropriate for distantly related taxa because the high substitution rate and site heterogeneity of mtDNA often lead to mutational saturation in protein-coding genes (see figure). This saturation biases the NI toward values <1 as species divergence increases. The smaller number of invertebrate mtDNA genomes currently available tends to force



more distant outgroup comparisons.

Bazin *et al.* rightfully emphasize the necessity of adequately testing for deviation from the neutral model for mtDNA, as with all loci. Further, the meta-analytical tools developed by Bazin *et al.* and others can help assess the time scale of selective sweeps relative to demographic events commonly considered by evolutionary biologists (e.g., effects of glaciation, high variance in reproductive success, and recent/incipient speciation). All genetic data come with complications, but we argue that it is inappropriate and unnecessary to dismiss the contribution that mtDNA sequence data—still one of the most powerful universal sources of genetic variation for nonmodel

◀ Neutrality indices simulated for a single nonrecombining 1 kb coding region. For each interspecific distance class, 100 coalescent simulations were performed comparing an ingroup taxon of $n = 10$ and expected within-species pairwise divergence of 2% to a single outgroup taxon. Simulations assume a transition:transversion ratio of 2 and a relative substitution rate of 2:1:20 for the first, second, and third codon positions, respectively. Thick horizontal bars indicate medians, and boxes include 50% of the distributions. The vertical line indicates the cutoff point used by Bazin *et al.* in their meta-analysis.

animals—can make to studies of conservation, taxonomy, and historical demography.

JOHN P. WARES,¹ PAUL H. BARBER,²

JEFFREY ROSS-IBARRA,¹ ERIK E. SOTKA,³

ROBERT J. TOONEN⁴

¹Department of Genetics, University of Georgia, Athens, GA 30606, USA. ²Department of Biology, Boston University, Boston, MA 02215, USA. ³Grice Marine Laboratory, College of Charleston, Charleston, SC 29412, USA. ⁴Hawai'i Institute of Marine Biology, University of Hawai'i at Manoa, Kane'ohe, HI 96744, USA.

References

1. S. Y. W. Ho, M. J. Phillips, A. Cooper, A. J. Drummond, *Mol. Biol. Evol.* **22**, 1561 (2005).
2. A. S. Gerber, R. Loggins, S. Kumar, T. E. Dowling, *Annu. Rev. Genet.* **35**, 539 (2001).
3. D. Rubinoff, B. S. Holland, *Syst. Biol.* **54**, 952 (2005).
4. R. Frankham, *Genet. Res.* **66**, 95 (1995).

Letters to the Editor

Letters (~300 words) discuss material published in *Science* in the previous 6 months or issues of general interest. They can be submitted through the Web (www.submit2science.org) or by regular mail (1200 New York Ave., NW, Washington, DC 20005, USA). Letters are not acknowledged upon receipt, nor are authors generally consulted before publication. Whether published in full or in part, letters are subject to editing for clarity and space.

Response

Berry and Wares *et al.* independently comment on our conclusion that mtDNA might not be a reliable marker of species population size and diversity. They introduce four arguments: (i) age of selective sweeps, (ii) census size versus effective size, (iii) distance to outgroup, and (iv) the usefulness of mtDNA despite selective sweeps.

1) Despite the fact that our study is based on comparisons between distantly related taxa, the selective sweeps we think have contributed to decreased mtDNA diversity in large populations must be recent ones, because they have

influenced the level of polymorphism observable within species.

2) We agree that effective population size can be very different from census population size, to an extent largely variable between species. Our analysis, however, recovers a positive relationship between nuclear genetic diversity and indicators of species abundance, indicating that effective and census populations sizes are correlated. The lack of relationship with mtDNA markers can therefore hardly be due to the census versus effective size problem, especially given the much larger data set analyzed.

3) It is true that very distant outgroups can bias the NI analysis because of saturation of the synonymous divergence (d_s), as neatly demonstrated by Wares' simulations. Our data set does not show strong variation of mitochondrial d_s across taxa: the average d_s is 0.262 in invertebrates versus 0.266 in vertebrates.

4) We do not mean to argue that mtDNA markers should be abandoned; there are many practical reasons why they can be useful. We strongly caution mtDNA users, however, that within-species mtDNA variations are likely to be influenced by natural

selection, especially in invertebrate species, where adaptation might be the rule. The age of the most recent mtDNA ancestor, in particular, should not be connected to any climatic, geologic, or biotic event unless confirmation is obtained from nuclear markers.

E. BAZIN, S. GLÉMIN, N. GALTIER
CNRS UMR 5171—Génome, Populations, Interactions, Adaptation—Université Montpellier 2, 34095 Montpellier Cedex 5, France.

TECHNICAL COMMENT ABSTRACT

Comment on "Population Size Does Not Influence Mitochondrial Genetic Diversity in Animals"

Connie J. Mulligan, Andrew Kitchen, Michael M. Miyamoto

Bazin *et al.* (Reports, 28 April 2006, p. 570) found no relationship between mitochondrial DNA (mtDNA) diversity and population size when comparing across large groups of animals. We show empirically that species with smaller populations, as represented by eutherian mammals, exhibit a positive correlation between mtDNA and allozyme variation, suggesting that mtDNA diversity may correlate with population size in these animals.

Full text at

www.sciencemag.org/cgi/content/full/314/5804/1390a